

## 38 Degrees Garden

This garden contains lush, large-leaved plants from the northern North Island that are more tropical-looking than many New Zealand natives. Many of the large-leaved trees are from Manawatawhi (Three Kings Islands) which is in the rohe of Ngati Kuri.

|   |   |                                  |
|---|---|----------------------------------|
| <i>Ackama rosifolia</i>                             | <i>Coprosma crassifolia x</i>               | <i>Freycinetia banksii</i>       |
| <i>Adiantum formosum</i>                            | <i>Coprosma repens</i>                      | <i>Fuscospora fusca</i>          |
| <i>Agathis australis</i>                            | <i>Coprosma macrocarpa</i>                  | <i>Gahnia lacera</i>             |
| <i>Alectryon excelsus subsp. grandis</i>            | <i>Coprosma neglecta</i>                    | <i>Gahnia pauciflora</i>         |
| <i>Alseuosmia banksii</i>                           | <i>Coprosma neglecta x</i>                  | <i>Gahnia sp</i>                 |
| <i>Alseuosmia macrophylla</i>                       | <i>Coprosma rhamnoides</i>                  | <i>Hebe 'Waitakere'</i>          |
| <i>Alseuosmia quercifolia</i>                       | <i>Coprosma parviflora</i>                  | <i>Helichrysum lanceolatum</i>   |
| <i>Astelia banksii</i>                              | <i>Coprosma propinqua</i>                   | <i>Hydrocotyle dissecta</i>      |
| <i>Astelia grandis</i>                              | <i>Coprosma rigida</i>                      | <i>Lastreopsis microsora</i>     |
| <i>Astelia hastata</i>                              | <i>Coprosma spathulata</i>                  | <i>Lepidosperma laterale</i>     |
| <i>Astelia solandri</i>                             | <i>Coprosma spathulata subsp. hikuruana</i> | <i>Libertia grandiflora</i>      |
| <i>Astelia trinervia</i>                            | <i>Cordyline australis</i>                  | <i>Libocedrus plumosa</i>        |
| <i>Beilschmiedia tarairi</i>                        | <i>Cordyline banksii</i>                    | <i>Litsea calicaris</i>          |
| <i>Blechnum discolor</i>                            | <i>Cordyline obtecta</i>                    | <i>Lophomyrtus bullata</i>       |
| <i>Blechnum durum</i>                               | <i>Corokia buddleioides</i>                 | <i>Lophomyrtus obcordata</i>     |
| <i>Brachyglottis perdicioides var. perdicioides</i> | <i>Corokia macrocarpa</i>                   | <i>Machaerina sinclairii</i>     |
| <i>Carex lambertiana</i>                            | <i>Dianella nigra</i>                       | <i>Melicope aff. ternata</i>     |
| <i>Coprosma aff. rhamnoides</i>                     | <i>Diplazium australe</i>                   | <i>Melicytus macrophyllus</i>    |
| <i>Coprosma crassifolia</i>                         | <i>Doodia sp</i>                            | <i>Melicytus novae-zelandiae</i> |
|   | <i>Elingamita johnsonii</i>                 | <i>Meryta sinclairii</i>         |
|   |   | <i>Metrosideros bartlettii</i>   |

## 38 Degrees Garden *continued* . .

|                                    |   |
|------------------------------------|---|
| <i>Metrosideros fulgens</i>        | <i>Pseudowintera insperata</i>              |
| <i>Metrosideros robusta</i>        | <i>Ptisana salicina</i>                     |
| <i>Metrosideros umbellata</i>      | <i>Raukaua anomalus</i>                     |
| <i>Microlaena avenacea</i>         | <i>Raukaua edgerleyi</i>                    |
| <i>Mida salicifolia</i>            | <i>Rhabdothamnus solandri</i>               |
| <i>Myrsine chathamica</i>          | <i>Rhopalostylis sapida</i> aff.            |
| <i>Nephrolepis cordifolia</i>      | 'Chatham Islands'                           |
| <i>Nestegis apetala</i> (Vahl)     | <i>Rhopalostylis sapida</i>                 |
| <i>Nestegis cunninghamii</i>       | <i>Streblus banksii</i>                     |
| <i>Nestegis lanceolata</i>         | <i>Streblus heterophyllus</i>               |
| <i>Olearia angulata</i>            | <i>Streblus smithii</i>                     |
| <i>Pennantia baylisiana</i>        | <i>Tecomanthe speciosa</i>                  |
| <i>Pennantia baylisiana</i> 'Otari | <i>Teucrium parvifolium</i>                 |
| Debut' x <i>Pennantia</i>          | <i>Toronia toru</i>                         |
| <i>corymbosa</i>                   | <i>Veronica adamsii</i>                     |
| <i>Pennantia corymbosa</i>         | <i>Veronica diosmifolia</i>                 |
| <i>Piper melchior</i>              | <i>Veronica flavida</i>                     |
| <i>Pittosporum umbellatum</i>      | <i>Veronica macrocarpa</i>                  |
| <i>Planchonella costata</i>        | <i>Veronica perbella</i>                    |
| <i>Podocarpus laetus</i>           | <i>Veronica punicea</i>                     |
| <i>Podocarpus totara</i>           | <i>Veronica rivalis</i>                     |
| <i>Prumnopitys taxifolia</i>       | <i>Veronica speciosa</i>                    |
| <i>Pseudopanax ferox</i>           | <i>Veronica stricta</i> var. <i>stricta</i> |
| <i>Pseudopanax gilliesii</i>       | <i>Vitex lucens</i>                         |
| <i>Pseudowintera axillaris</i>     |   |